



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document:

Wood, Matthew J ORCID: 0000-0003-0920-8396, Childs, Dylan Z, Davies, Alicia S, Hellgren, Olof, Cornwallis, Charlie K, Perrins, Chris M and Sheldon, Ben C (2013) The epidemiology underlying age-related avian malaria infection in a long-lived host: the mute swan *Cygnus olor*. *Journal of Avian Biology*, 44 (4). pp. 347-358. doi:10.1111/j.1600-048X.2013.00091.x

Official URL: <http://onlinelibrary.wiley.com/doi/10.1111/j.1600-048X.2013.00091.x/abstract>

DOI: <http://dx.doi.org/10.1111/j.1600-048X.2013.00091.x>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/569>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

This is a peer-reviewed, pre-print (final draft post-refereeing) version of the following published document:

Wood, Matthew J. and Childs, Dylan Z. and Davies, Alicia S. and Hellgren, Olof and Cornwallis, Charlie Kand and Perrins, Chris M. and Sheldon, Ben C. (2013). The epidemiology underlying age-related avian malaria infection in a long-lived host: the mute swan *Cygnus olor*. *Journal of Avian Biology*, 44 (4) 347-358.

Published in *Journal of Avian Biology*, and available online at:

<http://onlinelibrary.wiley.com/doi/10.1111/j.1600-048X.2013.00091.x/abstract>

We recommend you cite the published (post-print) version.

The URL for the published version is

<http://dx.doi.org/10.1111/j.1600-048X.2013.00091.x>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

1 The epidemiology underlying age-related avian malaria
2 infection in a long-lived host: the mute swan *Cygnus olor*

3
4 Matt J. Wood^{1,2,*}, Dylan Z. Childs³, Alicia S. Davies¹, Olof Hellgren^{1,4}, Charlie K.
5 Cornwallis^{1,4}, Chris M. Perrins¹ & Ben C. Sheldon¹

6
7 ¹ Edward Grey Institute, Department of Zoology, University of Oxford, Oxford,
8 OX1 3PS, United Kingdom

9 ² Current address: School of Natural & Social Sciences, University of
10 Gloucestershire, Cheltenham, GL50 4AZ, United Kingdom

11 ³ Department of Animal & Plant Sciences, University of Sheffield, Sheffield, S10
12 2TN, United Kingdom

13 ⁴ Current address: MEMEG, Department of Biology, Lund University, S-22362
14 Lund, Sweden

15
16 * Correspondence: mjwood@glos.ac.uk

17
18 **Keywords:**

19 Disease dynamics, Anatidae, Haemosporida

20

Abstract

Quantifying the factors that predict parasite outbreak and persistence is a major challenge for both applied and fundamental biology. Key to understanding parasite prevalence and disease outbreaks is determining at what age individuals show signs of infection, and whether or not they recover. Age-dependent patterns of the infection of a host population by parasites can indicate among-individual heterogeneities in their susceptibility to, or rate of recovery from, parasite infections. Here, we present a cross-sectional study of avian malaria in a long-lived bird species, the mute swan *Cygnus olor*, examining age-related patterns of parasite prevalence and modelling patterns of infection and recovery. 115 swans, ranging from one to nineteen years old, were screened for infection with *Plasmodium*, *Haemoproteus* and *Leucocytozoon* parasites. Infections with three cytochrome-*b* lineages of *Haemoproteus* were found (pooled prevalence 67%), namely WW1 (26%), which is common in passerine birds, and two new lineages closely related to WW1: MUTSW1 (25%) and MUTSW2 (16%). We found evidence for age-related infection in one lineage, MUTSW1. Catalytic models examining patterns of infection and recovery in the population suggested that infections in this population were not life-long – recovery of individuals was included in the best fitting models. These findings support the results of recent studies that suggest hosts can clear infections, although patterns of infection-related mortality in older birds remain to be studied in more detail.

Introduction

Parasite dynamics are the result of a complex interplay between exposure, transmission, disease-induced morbidity or death, immune-mediated recovery, and host life history (Hudson et al. 2002), each of which may be influenced by environmental or among-individual heterogeneities (Wilson 2002). Quantifying these basic processes allows us to understand the factors that govern parasite outbreaks and long-term persistence (Dobson and Foufopoulos 2001), and can reveal potentially important associations between biotic and abiotic components of the environment and parasite performance (Patz et al. 2000). Such studies are an essential prerequisite for the effective design of control and eradication programs, and provide a basis for delineating the fundamental drivers of parasite dynamics in space and time. In the absence of more detailed longitudinal study, the most commonly available data for inferring key epidemiological parameters is apparent prevalence, the proportion of animals in a population that test positive for an infection (Heisey et al. 2006). Such data are relatively easy to collect compared to logistically challenging longer term studies, and when coupled with measurements of the age of individuals provide a relatively straightforward means of comparing key epidemiological parameters among host populations or parasite lineages.

A number of cross-sectional studies have reported differences in the degree of parasitism in the wild, revealing variation in space or time (Altizer et al. 2006, Bensch and Åkesson 2003, Cosgrove et al. 2008, Loiseau et al. 2010, Wood et al. 2007). However, such studies do not always permit age-prevalence patterns to be examined in detail, either because age may not be known due to constraints in the marking of

individuals at birth, or because it may not be possible to infer age from morphological traits. Moreover, most studies of disease in wild bird populations have been conducted in study organisms that are short-lived relative to the scale of seasonal variation in transmission (Beadell et al. 2006, Ishtiaq et al. 2007, Ricklefs et al. 2005, Scheuerlein and Ricklefs 2004, Ventim et al. 2012). This highlights the need for more studies of parasite dynamics in longer-lived host species, which are poorly represented among avian studies (Bennett and Owens 2002)

In the absence of significant disease-induced mortality, the observed pattern of apparent prevalence is driven by rates of transmission and recovery. The direct measurement of transmission presents significant logistical challenges, requiring observation of contact rates and infection probabilities, both seldom directly observable outside the lab (for exceptions see Goeyvaerts et al. 2010, Kjaer et al. 2008). Insight into the transmission process is often gained indirectly, therefore, by estimating the force-of-infection (FOI), also known as the infection hazard, which is simply the per capita rate of infection of susceptible hosts. Though it does not reflect transmission per se (the density or frequency of infected individuals, and rates of transmission govern its magnitude), FOI is nonetheless an important metric for quantifying disease foothold in a population (Heisey et al. 2006, Long et al. 2010). In an endemic setting – one in which the disease is at, or near, dynamic equilibrium or at least changing slowly relative to the lifespan of the host – FOI can be estimated from age-prevalence data derived from cross-sectional sampling. Such data has long been used to estimate the force of infection in human populations using ‘catalytic’ models based on the proportional-hazards framework (Anderson and May 1985, Bundy et al. 1987, Farrington et al. 2001, Grenfell and

Anderson 1985, Keiding 1991, Keiding et al. 1996), and occasionally in a wildlife disease setting (Caley and Hone 2002, Heisey et al. 2006, Hudson and Dobson 1997, Woolhouse and Chandiwana 1992).

Avian malaria in long term study populations offers a useful model system for the study of ecological drivers of disease in wild populations. Avian malaria, *Plasmodium* and *Haemoproteus* spp. (*sensu* Pérez-Tris et al. 2005; see Valkiūnas et al. 2005 for a contrasting view), is a vector-borne disease transmitted by haematophagous Diptera. Avian *Plasmodium* is transmitted primarily by mosquitoes (Culicidae) and *Haemoproteus* by biting midges (Ceratopogonidae) and louse flies (Hippoboscidae) (Valkiūnas 2005). The development of sensitive and accurate molecular diagnosis techniques for avian malaria (Hellgren et al. 2004) has revealed a substantial and unexpected diversity of malaria lineages (Bensch et al. 2004), many showing marked variation between lineages in associations with biotic and abiotic factors (Cosgrove et al. 2008, Knowles et al. 2011, Wood et al. 2007).

In this paper we report the results of a cross-sectional survey of the prevalence of avian malaria parasites in a resident, colonial population of mute swans *Cygnus olor* in which the majority of individuals are of known age. Mute swans are relatively long-lived birds, with some individuals living beyond 20 years (Charmantier et al. 2006b, McCleery et al. 2002). There are relatively few reports of avian malaria in other swan species (Bennett et al. 1984, Ramey et al. 2012, Ricklefs and Fallon 2002, Valkiūnas 2005). We examine age-specific variation in the apparent prevalence of infection with specific lineages of avian malaria using non-parametric regression, and then construct catalytic

infection models to estimate FOI in the face of different assumptions governing recovery and re-infection of individuals.

Methods

Host sampling

Mute swans (family Anatidae) are large (7-14kg) waterbirds found in lakes, rivers and coastal areas in temperate and oceanic climates in Europe & Asia, feeding in shallow water on aquatic vegetation. In Western Europe, mute swans are usually territorial and nest in isolated pairs close to the water's edge (Cramp and Simmons 1983), but uniquely in the UK a colony of mute swans is located on the south coast of England at Abbotsbury Swannery (50°35'N, 2°30'W) in The Fleet, a 14km tidal lagoon. The colony has been in existence since at least the 1300s and its population dynamics studied since the late 1960s (Perrins and Ogilvie 1981). The breeding population has increased steadily to become relatively stable at approximately 130 breeding pairs (1990-2012: McCleery et al. 2002); C.M. Perrins unpublished data). Most swans in the colony are ringed as cygnets shortly after hatching and are natally philopatric – only 5% of breeding females are immigrants (Charmantier et al. 2006a) – so the age of most adult birds is known. The majority of breeding pairs nest near brackish pools, ditches and streams or in nearby *Phragmites* reedbeds. Supplemental feeding and the protection of vulnerable breeding pairs in pens are part of the long-term management of the population, but this intervention does not

appear to cause significant differences in population ecology between the Abbotsbury colony and territorial swans in other parts of the UK (Perrins and Ogilvie 1981).

In August 2008, 115 birds of known ages between 1 and 19 years were blood sampled by ulnar or tarsal venepuncture under UK Home Office licence. All were recruits to the colony (i.e. hatched as cygnets at Abbotsbury) to avoid the potential confounding effects of immigration. 92 of these birds were of known sex, based on the consensus from observations of sexually dimorphic bill knob size (Horrocks et al. 2006), sexual behaviour and cloacal examination (Swan Study Group 2005) made during the 2008 capture and any preceding captures or resightings.

Parasite screening

Samples were stored in SET buffer (0.015 M NaCl, 0.05 M Tris, 0.001 MEDTA, pH 8.0), and DNA extracted using a standard ammonium acetate protocol with the final product eluted in Qiagen AE buffer (Qiagen, Valencia CA, USA). DNA was quantified in 100x dilutions using pico-green dye and diluted to a final concentration of 25ng/μl. Extraction products were screened for the presence of *Leucocytozoon*, *Plasmodium* and *Haemoproteus* infections following the protocol by (Hellgren et al. 2004). 3μl of PCR product was run on 2% agarose gel containing ethidium bromide with a 1Kbp DNA ladder for each sample, testing for *Leucocytozoon* or *Haemoproteus/Plasmodium* lineages. Standard positives were used from previous malaria work in blue tits *Cyanistes caeruleus* (Knowles et al. 2010, Wood et al. 2007). Clear, strong bands were taken as positive and the absence of a band as negative for infection. Negative samples were rescreened to verify parasite absence. Positive sample PCR products were cleaned using a

Qiavac multiwell vacuum manifold.. To identify cytochrome-*b* lineages, BigDye (Applied Biosystems, Foster City, CA, USA) sequencing reactions were run with the forward primer for each lineage (F or FL). Sequences were edited and aligned in Sequencher 4.2 (GeneCodes Corp., Ann Arbor, MI, USA) using known, widespread malarial lineages as an alignment reference obtained from the MalAvi database (Bensch et al. 2009, accessed 18th February 2013). Novel lineages were identified by BLAST search against Genbank and then by comparison with all Haemosporidian lineages in the MalAvi database. Novel lineages were then named to indicate the bird host, using the five-letter species codes (as used by the British Trust for Ornithology), i.e. MUTSW for mute swan, suffixed by a number for each new lineage. Data on novel lineages were submitted to GenBank and MalAvi databases.

Phylogenetic analysis

To place the lineages found in this study in a phylogenetic context, we used the MalAvi database (Bensch et al. 2009) to identify (i) all lineages of *Haemoproteus*, *Plasmodium* and *Leucocytozoon* previously found in Anatidae, and (ii) the closest known relatives of the lineages found in this study. The latter were selected by constructing a neighbour-joining phylogeny – a Jukes-Cantor model as implemented in Geneious ver. R6 (Biomatters Ltd., Auckland, New Zealand: <http://www.geneious.com>) – including all available lineages to identify a well-defined clade of lineages in which the lineages found in this study occurred (for selection of clusters see Appendix 2). A Bayesian phylogeny was then constructed from the selected lineages using MrBayes (Huelsenbeck and Ronquist 2001) with a GTR-inv.gamma model allowing for 6 different gamma

categories. The total run length was 1000000 with trees sampled every 200th step. After discarding the first 10000 trees, the remaining trees were used to construct a consensus tree. The phylogeny was visualised using MEGA 5.0 (Kumar et al. 2008)

Statistical analysis

1. Age-related variation in prevalence

We used generalized additive models (GAM) to accommodate potentially non-linear relationships between parasite prevalence and age (Hudson et al. 2002, Wilson 2002). GAMs allow the expected value of the response to vary as a smooth function of a predictor (host age in this case) alongside conventional linear or categorical predictors and their interactions (Wood 2006). First, we analysed infection with each parasite lineage separately: starting models incorporated a smoothed function of age and host sex as model predictors, using binomial errors and a logit link. Beginning with the interaction term, which was introduced as separate smoothed age functions for each sex, predictors were eliminated from the model if removal resulted in a non-significant change in model deviance ($P > 0.05$), using likelihood ratio tests with penalised likelihoods and a backward stepwise procedure. Patterns of prevalence were visualized by calculating the predicted fitted response of each GAM of sample date on parasite infection: this approach applies the estimated model effects to a hypothetical range of sampling ages to calculate the fitted response and associated confidence estimates. GAMs were not forced through the origin (as would be appropriate for a disease without vertical transmission, i.e. zero prevalence at age zero), because birds in their first year of life (zero years) would, in fact, have been weeks or months old at the time of sampling (late summer) and therefore cannot be assumed to be free of avian malaria infection. To test directly for differences in

age-prevalence variation between parasite lineages, we used generalized additive mixed modelling (GAMM, Wood 2006). Each host individual was represented by three data per individual reflecting infection with each of three lineages, with individual identity fitted as a random effect and varying coefficient smoothing with respect to infection with each lineage. Once GAMM model selection was completed using maximum likelihood, the model was refitted using REML for extraction of parameter estimates. These analyses were conducted using the packages mgcv 1.7-13 and gamm4 0.1-5 in R 2.15.0 (R Core Team 2012). Means are presented ± 1 standard error.

2. Catalytic model

The observed age-prevalence data is multinomial, with four possible outcomes corresponding to individual infection status: uninfected ($Y = 0$), infected with lineage WW1 (see results) ($Y = 1$), infected with MUTSW1 ($Y = 2$), or infected with MUTSW2 ($Y = 3$). Two assumptions are made:

(i) That infections are avirulent, such that infection-induced mortality is negligible and can be ignored. This assumption was traditionally accepted by early studies of avian blood parasites (Bennett et al. 1988, Valkiūnas 2005). While a growing number of correlative studies have been equivocal on the virulence of avian malaria in stable host populations (Asghar et al. 2011, Atkinson et al. 2008, Lachish et al. 2011a), recent experimental studies have detected the detrimental effects of avian malaria in wild passerine bird populations (Knowles et al. 2010, Martinez-de la Puente et al. 2010, Marzal et al. 2005, Merino et al. 2000). In view of the equivocal evidence from avian malaria studies and the lack of evidence from long-lived birds in general (and our study

population in particular), we take the lack of virulence as a starting assumption for our catalytic models. We re-examine this assumption in the Discussion.

(ii) That the disease is endemic, such that disease incidence remains constant within the transmission period. Long-term studies of avian malaria in host populations are infrequent (Lachish et al. 2011b, Westerdahl et al. 2005), but generally do not support the existence of epidemic outbreaks of infection in stable host populations (Atkinson et al. 2008). Under these assumptions the age-prevalence data can be used to estimate the FOI, that is, the rate at which susceptible individuals acquire infection (Heisey et al. 2006). We can also estimate parameters describing what happens once individuals become infected for the first time: (i) infections are life-long and there is no recovery; (ii) individuals recover to become fully susceptible again; (iii) individuals recover and to possess lifelong immunity. Adopting the conventional language of epidemiology, we refer to these as the SI-, SIS-, and SIR-case, respectively. Since the FOI will differ among strains when either the prevalence or the transmission rates (or both) vary by strain, we consider two further possibilities for each scenario: the FOI is constant with respect to the strain; or the FOI varies by strain. Thus, we fit a total of six models to our age-prevalence data. In each, we assume that co-infection does not occur, an assumption justified as all sequence electropherograms were carefully examined for mixed infections and none was observed (Pérez-Tris and Bensch 2005). While we accept that these modelling assumptions place restrictions on the conclusions that may be drawn from our analyses, we believe that this modelling approach is justified by the utility of estimating FOI, which is often difficult to estimate in wild populations (McCallum et al. 2001), for example to enable comparison of FOI within and between studies.

247 In order to construct the likelihood under a particular model we need to calculate
 248 the probability that an individual has a given infection status (0, 1, 2, or 3) at each age, up
 249 to the maximum age observed. Because the FOI and recovery rates (if present) are
 250 constant with respect to age, the required likelihood can be calculated by iterating matrix
 251 projection model describing the transitions among infection states in successive ages. In
 252 the SI-case with strain varying FOI, the matrix projection model has the form

$$253 \quad \mathbf{p}_a = \Psi^a \mathbf{p}_0, \quad (1)$$

254 where \mathbf{p}_a denotes the distribution vector of states at age a , \mathbf{p}_0 is the initial distribution
 255 vector, and Ψ^a is the transition matrix for infection raised to the a^{th} power. Since
 256 individuals are uninfected at birth the initial distribution vector is simply $\mathbf{p}_0 = (1 \ 0 \ 0 \ 0)^T$.

257 The infection matrix Ψ has the form

$$258 \quad \Psi = \begin{pmatrix} \pi & 0 & 0 & 0 \\ (1-\pi)\theta_1 & 1 & 0 & 0 \\ (1-\pi)\theta_2 & 0 & 1 & 0 \\ (1-\pi)\theta_3 & 0 & 0 & 1 \end{pmatrix}, \quad (2)$$

259 where π is the probability an individual avoids infection over the course of a year and θ_i
 260 is the probability that an individual is infected by strain i . These probabilities are
 261 expressed in terms of the FOI for each strain, λ_i , such that $\pi = e^{-\lambda_1 - \lambda_2 - \lambda_3}$ and

262 $\theta_i = \frac{\lambda_i}{\lambda_1 + \lambda_2 + \lambda_3}$. The likelihood is then

$$263 \quad y_i \sim \text{Multinom}(1, \mathbf{p}_a), \quad (3)$$

264 where y_i denotes the status of individual i . Readers familiar with survival analysis will
 265 recognise that the FOI can also be estimated with a competing risks survival model,

where the “hazard” associated with each infection process is λ_i . However, we use the above formulation because it is easily extended to incorporate recovery.

In order to construct the likelihood for the SIS- and SIR-case we assume that the infection and recovery processes occur independently and sequentially each year. This approximation, which simplifies the modelling, is justified by the observation that insect vectors are absent over the winter, so that transmission necessarily occurs in late spring and summer. We also assume that within-year recovery during the spring/summer transmission period is negligible, and can be ignored; an assumption that may be justified on two counts: Firstly, previous studies of avian haemosporidia indicate that if a bird survives the initial acute phase of infection, usually occurring on being first exposed to infection as a juvenile, there follows a chronic phase of infection that persists for an extended period of time at low parasitaemia (Valkiūnas 2005, Zehtindjiev et al. 2008). Most individuals in this study were sampled as adults with no noticeable symptoms of infection, so we assume that these infections are in the chronic, stable, low intensity phase of infection and therefore unlikely to change infection status during the transmission period. Secondly, between-year repeatability of individual avian malaria infection is lower than within-year repeatability (Knowles et al. 2011). In the SIS-case with strain varying FOI, the matrix projection model used to construct the likelihood has the form

$$\mathbf{p}_a = (\Lambda \Psi)^a \mathbf{p}_0, \quad (4)$$

where \mathbf{p}_a , \mathbf{p}_0 and Ψ are defined as above and Λ denotes the transition matrix for recovery to the susceptible state. The recovery matrix Λ has the form

$$\Lambda = \begin{pmatrix} 1 & 1-\phi & 1-\phi & 1-\phi \\ 0 & \phi & 0 & 0 \\ 0 & 0 & \phi & 0 \\ 0 & 0 & 0 & \phi \end{pmatrix}, \quad (5)$$

where ϕ denotes the probability that an individual remains infected over autumn and winter. This probability is expressed in terms of the recovery rate, ρ , such that $\phi = e^{-\rho}$. The likelihood is then given by equation 3 above. In the SIR-case with strain varying FOI the matrix projection model used to construct the likelihood has the form

$$\mathbf{p}_a = (\Omega \Psi)^a \mathbf{p}_0, \quad (6)$$

where \mathbf{p}_a , \mathbf{p}_0 and Ψ are defined as above and Ω denotes the transition matrix for recovery to the immune state. The recovery matrix Ω has the form

$$\Omega = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \eta & 0 & 0 \\ 0 & 0 & \eta & 0 \\ 0 & 0 & 0 & \eta \end{pmatrix}, \quad (7)$$

where η denotes the probability that an individual remains infected over autumn and winter, which is expressed in terms of the recovery rate, γ , such that $\eta = e^{-\gamma}$. The likelihood is again given by equation 3 above, but with age specific distribution vector \mathbf{p}_a replaced with $\mathbf{p}'_a = (p'_{a,0} \ p_{a,1} \ p_{a,2} \ p_{a,3})$, where the $p_{a,i}$ are the elements of \mathbf{p}_a and $p_{a,0} = 1 - p_{a,1} - p_{a,2} - p_{a,3}$. This is because we do not (explicitly) track the immune class, yet the observed uninfected class includes both susceptible and recovered individuals.

The strain-independent FOI version of each model is obtained by simply setting $\lambda_1 = \lambda_2 = \lambda_3$ in infection matrix Ψ . Calculation of the likelihood and likelihood

maximisations were carried out with the R statistical programming language (R Core Team 2012). Univariate confidence intervals for model parameters were estimated from profile likelihoods. Models were compared using the small sample AIC_c, since the ratio of the number of observations to parameters in the highest dimensional models is small (Burnham and Anderson 2002).

3. Avian malaria infection and survival

To address one of the assumptions of catalytic modelling, that infection was avirulent, we scrutinised resighting and recapture data of swans sampled and screened for avian malaria infection in 2008. Every two years, approximately 99% of the population is captured during a ‘round-up’ of the swans at Abbotsbury; the identities of birds captured at round-ups in 2009 and 2011 was supplemented by data from individually colour-marked individuals resighted breeding in the colony in these years – typically around 50 breeding adults are not captured at each round-up (C.M. Perrins, unpublished data).

Therefore, we examine survival until 2009 and 2011 as two measures of mortality that may have been influenced by infection status at sampling in 2008. Included as factors in this analysis were infection status and individual age, the latter included both as age in years and categorised as young (0-9) or old (10-19 years) to examine the potential effects of infection-related mortality in later life. These data were analysed using a generalized additive model (Wood 2006) to examine the effect of parasite infection and age on survival until 2009 or 2011, with binomial errors and a logit link. Models were optimised by backward stepwise deletion: a predictor was deleted if its removal from the model made a non-significant change in model deviance (Analysis of deviance, $P > 0.05$).

Results

Avian malaria was diagnosed in 67.0% (77/115) of the mute swans sampled in this study. All infections belonged to the genus *Haemoproteus*, comprising the cytochrome-*b* lineage WW1 (prevalence 26.1%, 30/115) and two previously unreported lineages differing by one base pair difference in a 433bp cytochrome-*b* sequence, namely MUTSW1 (25.2%, 29/115: GenBank accession number GU319788) and MUTSW2 (15.7%, 18/115: GenBank accession number GU319789). All three lineages were very closely related to each other: MUTSW1 & 2 differed by just one nucleotide, and they in turn differed to WW1 by 2 and 3 substitutions respectively. The phylogenetic relationships between these lineages do not indicate a close connection with those previously found in Anatidae (Figure 1): these three lineages sit instead in a phylogenetic cluster containing *Haemoproteus* lineages previously found exclusively in passerine bird species (with the exception of lineage MEUND3, found in the budgerigar *Melopsittacus undulates*). For a full list of the hosts species in which the lineages shown in Figure 1 have been found, see Appendix 1. All lineages of *Plasmodium*, *Leucocytozoon* and *Haemoproteus* previously found in Anatidae are phylogenetically distant to the lineages found in this study.

Age-specific variation in prevalence

On examining the age-dependent pattern of infection of pooled *Haemoproteus* infections, infection appeared to rise steeply in older individuals, reaching a plateau of 80.6(±5.9)% prevalence at ten years of age: overall a significant age-prevalence relationship (GAM:

Analysis of deviance $\chi^2=14.6$, est.df=2.18, $P=0.020$; Figure 2a). Prevalence of infection with the three comprising lineages revealed varying patterns. WW1 and MUTSW2 showed no significant age-related variation ($P>0.05$; Figures 2b,d), however MUTSW1 infection showed significant age-related variation ($\chi^2=12.4$, est.df=3.21, $P=0.014$), increasing to a peak of $51.6(\pm 9.7)\%$ prevalence at approximately nine years of age before declining in older individuals (Figure 2c). In a direct test of age-dependent variation in infection, lineage identity did not have an overall significant effect on age-prevalence variation (GAMM: $\chi^2=3.67$, df=2, $P=0.16$), although a significant age:lineage interaction was detected for MUTSW1: the age-related pattern of infection for MUTSW1 was found to be significantly different to that of pooled infection with other *Haemoproteus* lineages ($\chi^2=12.0$, est.df=2.5, $P=0.0045$). Sex was not retained as a significant predictor of infection (GAM: $P>0.05$).

Epidemiological modelling

The relative performance of the six catalytic models is summarised in Table 1. The best model was the strain-independent force-of-infection (FOI) version of the susceptible-immune-recovered (SIR) model; this model predicts that the FOI does not vary among strains, and that individuals recover into a fully immune class. However, the AIC differences associated with the three remaining models that also include recovery processes were all less than 1, revealing very similar weights of evidence for these alternatives. The AIC difference of both models excluding the recovery were greater than 10, indicating that models excluding a recovery process were (relatively) very poor approximating models for the age-prevalence data. Taken together, these results provide

strong evidence against the possibility that *Haemoproteus* infections are life-long in the Abbotsbury mute swan population. However, our analysis was unable to resolve the nature of the recovery process (i.e. no significant preference between susceptible-immune-susceptible (SIS) and SI-recovered (SIR) models), and was not able to establish unequivocally whether the FOI varies among strains.

The maximum likelihood estimate of the per-strain FOI under the best model (SIR) is 0.10 (0.07-0.15, 95% CI), which implies that the annual probability of infection by any strain is 0.26 (0.18-0.36, 95% CI), whereas the estimated per-strain FOI under the strain-independent FOI model (capturing recovery back into a susceptible class: SIS) was very similar to that of the best model (0.12, 0.07-0.25 95% CI), implying a similar annual probability of infection by any strain (0.30, 0.20-0.53 95% CI). Although the predicted recovery rate under this alternative model was relatively higher than that of the best model, this rate is still very low (0.076, 0.023-0.19 95% CI) and corresponds to an annual recovery probability of only 0.08 (0.02-0.18 95% CI).

The predicted age-prevalence curves under each of the fitted models are summarised in Figure 3. Figure 3a shows the predicted relationship under the models assuming strain-independent FOI. It is clear that both the SIR and SIS models yield very similar relationships, which explains why we were unable to resolve differences among the two types of model. The remaining three figures (3b-d) summarise the predictions derived from the models allowing between-strain variation in the FOI. Though among strain differences in the predicted age-prevalence relationships can be detected ‘by eye’, this variation is very low (Table 1).

3. Avian malaria infection and survival

Infection with avian malaria at capture in 2008, either as pooled or as individual *Haemoproteus* lineages, was not associated with subsequent survival until August 2009, and age was not a contributory factor whether incorporated as a potentially non-linear effect or categorised as ‘young’ (0-9 years) or ‘old’ (10-19 years) (GAM: $P>0.3$). Age-dependent mortality was detected in data on survival until 2011 (as expected, older birds were less likely to survive: $\chi^2=8.18$, est.df=1.61, $P=0.017$), but this effect was not related to infection, either as pooled or lineage-specific *Haemoproteus* infections (GAM: age:infection interactions, $P>0.5$). Given the few old birds in our sample (40/115, 34.8%), we interpret the results of these basic analyses of survival with caution.

Discussion

We found that 67% of the mute swan population at Abbotsbury was infected with avian malaria, with one of three cytochrome-*b* lineages of *Haemoproteus*: namely WW1 (prevalence 26%) and two novel lineages not previously reported in previous studies, named MUTSW1 (25%) and MUTSW2 (15%). We found evidence for different age-prevalence patterns between lineages, with only MUTSW1 showing significant variation in prevalence with age, although this was not reflected in varying epidemiological parameters between lineages. Catalytic modelling found most support for models of age-dependent prevalence models including recovery from infection, rejecting models of life-long infection by a considerable margin.

This study suggests that the convex age-prevalence curve of *Haemoproteus* lineage MUTSW1 in mute swans at Abbotsbury (Figure 2c) may be due to hosts recovering from infection. These results contrast to some extent with previous studies, mainly based on longitudinal studies of short-lived captive birds (Atkinson et al. 2008, Palinauskas et al. 2008, Valkiūnas 2005), which observed an initial critical phase of infection that the host may or may not survive, followed by a decrease in the number of parasites in the bloodstream to a low, stable level that continues for an extended period of time, perhaps the remaining life time of the host (Valkiūnas 2005). Chronic infections are typically of low parasitaemia, but above the detection threshold of the molecular diagnosis techniques applied in this study (Knowles et al. 2010, Palinauskas et al. 2008), so it is unlikely that the loss of infection we report here is a result of a reduction of parasitaemia to undetectable levels. Mute swans live much longer than bird hosts examined in previous studies examining age-related patterns of malaria infection, approximately 10 years on average (McCleery et al. 2002), so host age-related parasite dynamics may be different in short and long lived birds: the fall in MUTSW1 prevalence in older mute swans may be a result of mechanisms that are not apparent in relatively short lived hosts. Acquired immunity (Anderson and May 1985, Crombie and Anderson 1985, Dobson et al. 1990, Woolhouse et al. 1991), and age-related behavioural variation in exposure to infection (Altizer et al. 2003, Halvorsen 1986) may contribute to the convex age-prevalence curve of MUTSW1, but further study would be required to identify the mechanisms involved. Parasite-mediated viability selection may also result in a convex age-infection pattern: higher mortality of infected young individuals will remove them from the population to result in a lower prevalence of disease in older

individuals (Sol et al. 2003, van Oers et al. 2010). Age-specific patterns of infection may simply remain as echoes of past epidemics, with some age cohorts retaining chronic disease infection acquired in previous outbreaks of a disease no longer transmitted in a population (Long et al. 2010). Furthermore, if senescence results in higher parasite-induced mortality in older individuals, perhaps mediated by a deteriorating immune system in older individuals (Vleck et al. 2007, Lavoie 2005), then older infected individuals will be removed from the population with a subsequent decline in the prevalence and intensity of infection. Although we found no evidence for an interaction between infection and mortality in this study, this may have been masked by our cross-sectional ‘snapshot’ sampling – examining this potential mechanism to explain declining prevalence with age would be an important goal for future studies, particularly to [scrutinise infection dynamics and mortality in older individuals](#). Further work would be necessary to resolve the potential mechanisms underlying age-specific variation in avian malaria infection in this population, and in wild populations more generally, involving detailed longitudinal studies of marked individuals combined with extensive infection screening (Lachish et al. 2011b, Westerdahl et al. 2005, Atkinson and Samuel 2010, van Oers et al. 2010), which would enable (i) the estimation of transitions between infection states (Atkinson and Samuel 2010, Faustino et al. 2004, Jennelle et al. 2007, Lachish et al. 2011a, Senar and Conroy 2004), (ii) a more detailed monitoring of infection-dependent mortality that may contribute to age-prevalence patterns, and (iii) the examination of the widely indicated importance of transmission in early life (Cosgrove et al. 2008, Valkiūnas 2005, Hasselquist et al. 2007). The benefits of molecular diagnosis in such studies are clear, but it would be important to include the preparation of blood films

to complement molecular diagnoses: Haemosporidian parasites might sometimes infect non-natural hosts without completing their lifecycle to the infective gametocyte stage, perhaps allowing degraded parasite DNA to be detected in the blood and thus the erroneous conclusion that the presence of a DNA lineage is evidence of a competent host (see Olias et al 2011 for an example of abortive development). Although it is unlikely that the three mute swan lineages detected in this study are examples of abortive development (high prevalence of infection, close relationship between lineages), the collection of data for microscopy remains important in an age of molecular diagnostics.

Assumptions of the epidemiology of avian malaria used in the construction of catalytic models should be realistic for their results to be reliable. Avian malaria parasites were assumed to be avirulent; traditionally accepted from reports of mortality confined to outbreaks in poultry (Atkinson et al. 2008, Valkiūnas 2005) but contradicted by recent experimental studies of wild bird populations that detect more subtle negative effects of infection on reproductive effort and success (Knowles et al. 2010, Marzal et al. 2005, Merino et al. 2000) and survival (Martinez-de la Puente et al. 2010). While a general pattern exists for avian malaria infection to be negatively correlated with reproductive effort (Knowles et al. 2009), detecting the consequences of infection as host mortality have proved more elusive. All the infections detected in the current study were *Haemoproteus*, generally accepted to be less virulent than *Plasmodium* or *Leucocytozoon* haemosporidian parasites in stable host populations, in contrast with studies of the devastating effects of avian *Plasmodia* introduced to naïve host populations (Atkinson and Samuel 2010). Recapture/resighting of the swans in this study since sampling in 2008 revealed no significant effect of infection on the probability of survival. Although an

experimental approach would be preferable, we do not detect any obvious consequences of *Haemoproteus* infection for survival in this mute swan colony, and suggest that our assumption of avirulent *Haemoproteus* infections is a reasonable working hypothesis with which to estimate the force-of-infection (FOI), a valuable epidemiological parameter.

Few studies have examined water birds for avian malaria infection using modern molecular diagnostic techniques, although two studies report avian malaria in the Tundra swan *Cygnus columbianus* in North America (Ricklefs and Fallon 2002, Ramey et al. 2012). Considering the distribution of the avian malaria lineages previously reported in swans in this and other studies (Bensch et al. 2009) reveals that the *Haemoproteus* lineage WW1 has previously been found in ten bird species, all of which are passerine birds from the Western Palaearctic or (with the exception of the paddyfield warbler *Acrocephalus agricola*) from the Palaearctic-African migratory flyway (Bensch and Åkesson 2003, Hellgren et al. 2007, Krizanaskiene et al. 2006, Ventim et al. 2012, Wood et al. 2007). WW1 is the closest known relative to the other two lineages found in this study, so if the lineages MUTSW1/2 are exclusive to mute swans it is possible that a host shift has occurred whereby WW1 shifted into Anatidae and subsequently diversified into MUTSW1 & 2 (Figure 1). Avian malaria lineages of the genus *Haemoproteus* are known to be more host-specific than *Plasmodium* (Beadell et al. 2004), but WW1's appearance in mute swans would appear to be an exception to this reported pattern. It is clear from the phylogenetic relationship of *Haemoproteus* lineages in this study that infection is not always related to the evolutionary history of that host: hosts of relatively distant shared ancestry may share highly similar blood parasites (Figure 1). *Leucocytozoon* infection is

common in wildfowl, being a confirmed cause of mortality in young Anatidae (ducks, geese and swans) including mute swans (Mörner and Wahlström 1983, Valkiūnas 2005), yet no such infections were found in this mute swan colony, perhaps due to the inability of blackfly vectors to tolerate the tidally variable salinity of The Fleet lagoon at the Abbotsbury Swannery (Williams and Williams 1998). The lack of malaria screening in water birds making it difficult to draw comparisons with other, more intensively surveyed taxa (mostly Passeriformes), generally smaller birds that can be conveniently sampled by mist-netting or while breeding in artificial nest boxes.

In conclusion, this study found a high prevalence of avian malaria infection in a mute swan colony comprised of three lineages of *Haemoproteus*, two of which were novel. Age-dependent variation in infection was found for just one of these lineages. Catalytic modelling provided strong evidence for recovery from *Haemoproteus* infections. The Abbotsbury mute swan colony would be a useful model system for the further study of pathogen outbreaks and persistence: the long-term study of individually-marked populations has brought considerable benefits to the fundamental understanding of ecology and evolution (Clutton-Brock and Sheldon 2010). The integration of systematic diagnosis of disease in wild populations has the potential to stimulate advances in the ecology and evolution of infectious disease.

Acknowledgements

We are very grateful to Mrs. C. Townshend, the owner of Abbotsbury Swannery, for permission to study the swans and to all those who assisted with population monitoring,

in particular Dave Wheeler and his staff at Abbotsbury. Ruth Cromie and Debbie Pain (Wildfowl and Wetlands Trust), Alasdair Dawson (CEH Edinburgh), and Miriam Liedvogel (Edward Grey Institute) assisted with blood sampling. Valuable support was provided to ASD by Sarah Knowles and Ricardo Alves in the lab. This work was supported by a Natural Environment Research Council standard research grant to BCS, a Natural Environment Research Council standard research fellowship to DZC, and by the Biotechnology and Biological Sciences Research Council's funding of The University of Oxford's M.Sc. course in Integrated Biosciences.

Figure Legends

Figure 1.

A Bayesian phylogram of lineages found in this study (indicated by arrows), with lineages previously found in hosts belonging to the Anatidae (+) and a selection of closely-related lineages (see Appendix 1). 'P' marks lineages previously found in passerine birds, 'U' those found only in the budgerigar *Melopsittacus undulatus*. All lineages belonging to the genus *Haemoproteus* marked with a P have been found previously only in passerine birds (for full list of hosts see Appendix 1). Numbers on branches represent posterior probabilities.


Figure 2.


Age-specific variation in avian malaria infection. (a) Pooled infections, all of which were *Haemoproteus* spp., and (b-d) showing variation in three cytochrome-*b* lineages of *Haemoproteus*. The size of points reflects the sample size, bars indicate standard error.

Figure 3.

Predicted age-prevalence patterns of mute swan malaria from catalytic models. Results of models for (a) pooled *Haemoproteus* infections and (b-d) infection with three cytochrome-*b* *Haemoproteus* lineages. Solid lines indicate predicted age-prevalence curves for the fitted SIR-case without strain-varying force of infection; dashed lines show the fitted SIS-case without strain-varying force of infection; and dotted lines show the fitted SI-case without strain-varying force of infection. Point size reflects sample size.

Figure 1.








Table 1.

Performance of the six catalytic models fitted to age-prevalence data of *Haemoproteus* infection in swans at Abbotsbury. Δ_i values in bold type indicate the models that were within 2 AIC units of the best model.

Model	Log Likelihood	Small sample AIC (AIC_c)	Number of parameters	AIC differences (Δ_i)
<i>SI</i>	-155.3	316.8	3	11.7
<i>SI</i> _{$\lambda_1 = \lambda_2 = \lambda_3$}	-157.1	316.3	1	11.2
<i>SIS</i>	-148.8	305.9	4	0.8
<i>SIS</i> _{$\lambda_1 = \lambda_2 = \lambda_3$}	-150.6	305.3	2	0.2
<i>SIR</i>	-148.7	305.7	4	0.6
<i>SIR</i> _{$\lambda_1 = \lambda_2 = \lambda_3$}	-150.5	305.1	2	0.0

582 References

- 583 Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M. and Rohani, P. 2006.
584 Seasonality and the dynamics of infectious diseases. – *Ecol. Lett.* 9: 467-484.
- 585 Altizer, S., Nunn, C. L., Thrall, P. H., Gittleman, J. L., Antonovics, J., Cunningham, A.
586 A., Dobson, A. P., Ezenwa, V., Jones, K. E., Pedersen, A. B., Poss, M. and
587 Pulliam, J. R. C. 2003. Social organization and parasite risk in mammals:
588 Integrating theory and empirical studies. – *Ann. Rev. Ecol. Evol. Syst.* 34: 517-
589 547.
- 590 Anderson, R. M. and May, R. M. 1985. Age-related changes in the rate of disease
591 transmissions - implications for the design of vaccination programs. – *J. Hyg.* 94:
592 365-436.
- 593 Asghar, M., Hasselquist, D. and Bensch, S. 2011. Are chronic avian haemosporidian
594 infections costly in wild birds? – *J. Avian Biol.* 42: 530-537.
- 595 Atkinson, C. T. and Samuel, M. D. 2010. Avian malaria *Plasmodium relictum* in native
596 Hawaiian forest birds: epizootiology and demographic impacts on 'apapane
597 *Himatione sanguinea*. – *J. Avian Biol.* 41: 357-366.
- 598 Atkinson, C. T., Thomas, N. J. and Hunter, D. B. 2008. Parasitic diseases of wild birds. –
599 Wiley-Blackwell, Ames, Iowa.
- 600 Beadell, J. S., Gering, E., Austin, J., Dumbacher, J. P., Peirce, M. A., Pratt, T. K.,
601 Atkinson, C. T. and Fleischer, R. C. 2004. Prevalence and differential host-
602 specificity of two avian blood parasite genera in the Australo-Papuan region. –
603 *Mol. Ecol.* 13: 3829-3844.
- 604 Beadell, J. S., Ishtiaq, F., Covas, R., Melo, M., Warren, B. H., Atkinson, C. T., Bensch,
605 S., Graves, G. R., Jhala, Y. V., Peirce, M. A., Rahmani, A. R., Fonseca, D. M. and
606 Fleischer, R. C. 2006. Global phylogeographic limits of Hawaii's avian malaria. –
607 *P. Roy. Soc. B - Biol. Sci.* 273: 2935-2944.
- 608 Bennett, G. F., Caines, J. R. and Bishop, M. A. 1988. Influence of blood parasites on the
609 body mass of passeriform birds. – *J. Wildl. Dis.* 24: 339-343.
- 610 Bennett, G. F., Turner, B. and Whiteway, M. 1984. Avian Haemoproteidae. XVIII:
611 *Haemoproteus greineri*, a new species of haemoproteid from the waterfowl family
612 Anatidae. – *Can. J. Zool.* 62: 2290-2292.
- 613 Bennett, P. M. and Owens, I. P. F. 2002. Evolutionary ecology of birds: life histories,
614 mating systems, and extinction. – Oxford University Press, Oxford.
- 615 Bensch, S. and Åkesson, A. 2003. Temporal and spatial variation of hematozoans in
616 Scandinavian willow warblers. – *J. Parasitol.* 89: 388-391.
- 617 Bensch, S., Hellgren, O. and Pérez-Tris, J. 2009. MalAvi: a public database of malaria
618 parasites and related haemosporidians in avian hosts based on mitochondrial
619 cytochrome b lineages. – *Mol. Ecol. Resour.* 9: 1353-1358.
- 620 Bensch, S., Pérez-Tris, J., Waldenström, J. and Hellgren, O. 2004. Linkage between
621 nuclear and mitochondrial DNA sequences in avian malaria parasites: Multiple
622 cases of cryptic speciation? – *Evol.* 58: 1617-1621.
- 623 Bundy, D. A. P., Cooper, E. S., Thompson, D. E., Anderson, R. M. and Didier, J. M.
624 1987. Age-related prevalence and intensity of *Trichuris trichuria* infection in a St.
625 Lucian community. – *Trans. Roy. Soc. Trop. Med.* H 81: 85-94.

- Burnham, K. P. and Anderson, D. R. 2002. Model Selection and Multi-Model Inference: A Practical Information-Theoretic Approach. – Springer, New York.
- Caley, P. and Hone, J. 2002. Estimating the force of infection; *Mycobacterium bovis* infection in feral ferrets *Mustela furo* in New Zealand. – J. Anim. Ecol. 71: 44-54.
- Charmantier, A., Perrins, C., McCleery, R. H. and Sheldon, B. C. 2006a. Age-dependent genetic variance in a life-history trait in the mute swan. – P. Roy. Soc. B - Biol. Sci. 273: 225-232.
- Charmantier, A., Perrins, C., McCleery, R. H. and Sheldon, B. C. 2006b. Quantitative genetics of age at reproduction in wild swans: Support for antagonistic pleiotropy models of senescence. – P. Natl. Acad. Sci. USA 103: 6587-6592.
- Clutton-Brock, T. and Sheldon, B. C. 2010. Individuals and populations: the role of long-term, individual-based studies of animals in ecology and evolutionary biology. – Trends Ecol. Evol. 25: 562-573.
- Cosgrove, C. L., Wood, M. J., Day, K. P. and Sheldon, B. C. 2008. Seasonal variation in *Plasmodium* prevalence in a population of blue tits *Cyanistes caeruleus*. – J. Anim. Ecol. 77: 540-548.
- Cramp, S. and Simmons, K. E. L. (eds.). 1983. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. - Oxford University Press, Oxford.
- Crombie, J. A. and Anderson, R. M. 1985. Population dynamics of *Schistosoma mansoni* in mice repeatedly exposed to infection. – Nature 315: 491-493.
- Dobson, A. and Foufopoulos, J. 2001. Emerging infectious pathogens of wildlife. – Philos. Trans. R. Soc. Lond. B Biol. Sci. 356: 1001-1012.
- Dobson, R. J., Waller, P. J. and Donald, A. D. 1990. Population dynamics of *Trichostrongylus colubriformis* in sheep - the effect of host age on the establishment of infective larvae. – Int. J. Parasit. 20: 353-357.
- Farrington, C. P., Kanaan, M. N. and Gay, N. J. 2001. Estimation of the basic reproduction number for infectious diseases from age-stratified serological survey data. – J. Roy. Stat. Soc. C - App. 50: 251-283.
- Faustino, C. R., Jennelle, C. S., Connolly, V., Davis, A. K., Swarthout, E. C., Dhondt, A. A. and Cooch, E. G. 2004. *Mycoplasma gallisepticum* infection dynamics in a house finch population: seasonal variation in survival, encounter and transmission rate. – J. Anim. Ecol. 73: 651-669.
- Goeyvaerts, N., Hens, N., Ogunjimi, B., Aerts, M., Shkedy, Z., Damme, P. V. and Beutels, P. 2010. Estimating infectious disease parameters from data on social contacts and serological status. – J. Roy. Stat. Soc. C - App. 59: 255-277.
- Grenfell, B. T. and Anderson, R. M. 1985. The estimation of age-related rates of infection from case notifications and serological data. – J. Hyg. 95: 419-436.
- Hall, T. A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. – Nucl. Acid. S. 41: 95-98.
- Halvorsen, O. 1986. On the relationship between social status of host and risk of parasitic infection. – Oikos 47: 71-74.
- Hasselquist, D., Ostman, O., Waldenström, J. and Bensch, S. 2007. Temporal patterns of occurrence and transmission of the blood parasite *Haemoproteus payevskyi* in the great reed warbler *Acrocephalus arundinaceus*. – J. Ornithol. 148: 401-409.

671 Heisey, D. M., Joly, D. O. and Messier, F. 2006. The fitting of general force-of-infection
672 models to wildlife disease prevalence data. – *Ecology* 87: 2356-2365.

673 Hellgren, O., Waldenström, J. and Bensch, S. 2004. A new PCR assay for simultaneous
674 studies of *Leucocytozoon*, *Plasmodium*, and *Haemoproteus* from avian blood. – *J.*
675 *Parasitol.* 90: 797-802.

676 Hellgren, O., Waldenström, J., Pérez-Tris, J., Szöllösi, E., Hasselquist, D.,
677 Krizanauskienė, A., Ottosson, U. and Bensch, S. 2007. Detecting shifts of
678 transmission areas in avian blood parasites - a phylogenetic approach. – *Mol.*
679 *Ecol.* 16: 1281-1290.

680 Horrocks, N., Sheldon, B., Perrins, C. and Charmantier, A. 2006. Bill knob size in the
681 mute swan: Evidence for a sexually selected trait? – *J. Ornithol.* 147: 183-184.

682 Hudson, P. J. and Dobson, A. P. 1997. Transmission dynamics and host-parasite
683 interactions of *Trichostrongylus tenuis* in red grouse (*Lagopus lagopus scoticus*).
684 – *J. Parasitol.* 83: 194-202.

685 Hudson, P. J., Rizzoli, A., Grenfell, B. T. and Dobson, A. P. (eds.). 2002. The Ecology of
686 Wildlife Diseases. - Oxford University Press, Oxford, U.K.,

687 Huelsenbeck, J. P. and Ronquist, F. 2001. MRBAYES: Bayesian inference of
688 phylogenetic trees. – 17: 754-755.

689 Ishtiaq, F., Gering, E., Rappole, J. H., Rahmani, A. R., Jhala, Y. V., Dove, C. J.,
690 Milensky, C., Olson, S. L., Peirce, M. A. and Fleischer, R. C. 2007. Prevalence
691 and diversity of avian hematozoan parasites in Asia: a regional survey. – *J. Wildl.*
692 *Dis.* 43: 382-398.

693 Jennelle, C. S., Cooch, E. G., Conroy, M. J. and Senar, J. C. 2007. State-specific
694 detection probabilities and disease prevalence. – *Ecol. Appl.* 17: 154-167.

695 Keiding, N. 1991. Age-specific incidence and prevalence - a statistical perspective. – *J.*
696 *Roy. Stat. Soc. A - Sta.* 154: 371-412.

697 Keiding, N., Begtrup, K., Scheike, T. H. and Hasibeder, G. 1996. Estimation from
698 current-status data in continuous time. – *Lifetime Data Anal.* 2: 119-129.

699 Kjaer, L. J., Schaubert, E. M. and Nielsen, C. K. 2008. Spatial and temporal analysis of
700 contact rates in female white-tailed deer. – *J. Wildl. Manage.* 72: 1819-1825.

701 Knowles, S. C. L., Nakagawa, S. and Sheldon, B. C. 2009. Elevated reproductive effort
702 increases blood parasitaemia and decreases immune function in birds: a meta-
703 regression approach. – *Funct. Ecol.* 23: 405-415.

704 Knowles, S. C. L., Palinauskas, V. and Sheldon, B. C. 2010. Chronic malaria infections
705 increase family inequalities and reduce parental fitness: experimental evidence
706 from a wild bird population. – *J. Evol. Biol.* 23: 557-569.

707 Knowles, S. C. L., Wood, M. J., Alves, R., Wilkin, T. A., Bensch, S. and Sheldon, B. C.
708 2011. Molecular epidemiology of malaria prevalence and parasitaemia in a wild
709 bird population. – *Mol. Ecol.* 20: 1062-1076.

710 Krizanaskienė, A., Hellgren, O., Kosarev, V., Sokolov, L., Bensch, S. and Valkiūnas, G.
711 2006. Variation in host specificity between species of avian hemosporidian
712 parasites: Evidence from parasite morphology and cytochrome B gene sequences.
713 – *J. Parasitol.* 92: 1319-1324.

714 Kumar, S., Nei, M., Dudley, J. and Tamura, K. 2008. MEGA: A biologist-centric
715 software for evolutionary analysis of DNA and protein sequences. – *Brief.*
716 *Bioinform.* 9: 299-306.

- Lachish, S., Knowles, S. C. L., Alves, R., Wood, M. J. and Sheldon, B. C. 2011a. Fitness effects of endemic malaria infections in a wild bird population: the importance of ecological structure. – *J. Anim. Ecol.* 80: 1196-1206.
- Lachish, S., Knowles, S. C. L., Alves, R., Wood, M. J. and Sheldon, B. C. 2011b. Infection dynamics of endemic malaria in a wild bird population: parasite species-dependent drivers of spatial and temporal variation in transmission rates. – *J. Anim. Ecol.* 80: 1207-1216.
- Lavoie, E. T. 2005. Avian immunosenescence. – *Age* 27: 281-285.
- Loiseau, C., Iezhova, T., Valkiūnas, G., Chasar, A., Hutchinson, A., Buermann, W., Smith, T. B. and Sehgal, R. N. M. 2010. Spatial variation of haemosporidian parasite infection in African rainforest bird species. – *J. Parasitol.* 96: 21-29.
- Long, G. H., Sinha, D., Read, A. F., Pritt, S., Kline, B., Harvill, E. T., Hudson, P. J. and Bjornstad, O. N. 2010. Identifying the age cohort responsible for transmission in a natural outbreak of *Bordetella bronchiseptica*. – *PLoS Pathog.* 6: e1001224.
- Martinez-de la Puente, J., Merino, S., Tomas, G., Moreno, J., Morales, J., Lobato, E., Garcia-Fraile, S. and Jorge Belda, E. 2010. The blood parasite *Haemoproteus* reduces survival in a wild bird: a medication experiment. – *Biol. Lett.* 6: 663-665.
- Marzal, A., de Lope, F., Navarro, C. and Moller, A. P. 2005. Malarial parasites decrease reproductive success: an experimental study in a passerine bird. – *Oecologia* 142: 541-545.
- McCallum, H., Barlow, N. and Hone, J. 2001. How should pathogen transmission be modelled? – *Trends Ecol. Evol.* 16: 295-300.
- McCleery, R. H., Perrins, C., Wheeler, D. and Groves, S. 2002. Population structure, survival rates and productivity of mute swans breeding in a colony at Abbotsbury, Dorset, England. – *Waterbirds* 25: 192-201.
- Merino, S., Moreno, J., Sanz, J. J. and Arriero, E. 2000. Are avian blood parasites pathogenic in the wild? A medication experiment in blue tits (*Parus caeruleus*). – *P. Roy. Soc. B - Biol. Sci.* 267: 2507-2510.
- Mörner, T. and Wahlström, K. 1983. Infektioner med blodparasiten *Leucocytozoon simondi* – en vanlig dödsorsak hos knölsvanungar *Cygnus olor*. – *Vår Fågelvärld* 42: 389-394.
- Olias, P., Wegelin, M., Zenker, W., Freter, S., Gruber, A. D. and Klopffleisch, R. 2011. Avian Malaria Deaths in Parrots, Europe. – *Emerg. Infect. Dis* 17: 950-952.
- Palinauskas, V., Valkiūnas, G., Bolshakov, C. V. and Bensch, S. 2008. *Plasmodium relictum* (lineage P-SGS1): Effects on experimentally infected passerine birds. – *Exp. Parasitol.* 120: 372-380.
- Patz, J. A., Graczyk, T. K., Geller, N. and Vittor, A. Y. 2000. Effects of environmental change on emerging parasitic diseases. – *Int. J. Parasit.* 30: 1395-1405.
- Pérez-Tris, J. and Bensch, S. 2005. Diagnosing genetically diverse avian malarial infections using mixed-sequence analysis and TA-cloning. – *Parasitol.* 131: 15-23.
- Pérez-Tris, J., Hasselquist, D., Hellgren, O., Krizanauskiene, A., Waldenström, J. and Bensch, S. 2005. What are malaria parasites? – *Trends Parasitol.* 21: 209-211.
- Perrins, C. M. and Ogilvie, M. A. 1981. A study of the Abbotsbury Mute Swans. – *Wildfowl* 32: 35-47.

- R Core Team 2012. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ramey, A. M., Ely, C. R., Schmutz, J. A., Pearce, J. M. and Heard, D. J. 2012. Molecular detection of hematozoa infections in tundra swans relative to migration patterns and ecological conditions at breeding grounds. – PLoS One 7: e45789-e45789.
- Ricklefs, R. E. and Fallon, S. M. 2002. Diversification and host switching in avian malaria parasites. – P. Roy. Soc. B - Biol. Sci. 269: 885-892.
- Ricklefs, R. E., Swanson, B. L., Fallon, S. M., Martinez-Abrain, A., Scheuerlein, A., Gray, J. and Latta, S. C. 2005. Community relationships of avian malaria parasites in southern Missouri. – Ecol. Monogr. 75: 543-559.
- Scheuerlein, A. and Ricklefs, R. E. 2004. Prevalence of blood parasites in European passeriform birds. – P. Roy. Soc. B - Biol. Sci. 271: 1363-1370.
- Senar, J. C. and Conroy, M. J. 2004. Multi-state analysis of the impacts of avian pox on a population of Serins (*Serinus serinus*): the importance of estimating recapture rates. – Anim. Biodiv. Conserv. 27: 133-146.
- Sol, D., Jovani, R. and Torres, J. 2003. Parasite mediated mortality and host immune response explain age-related differences in blood parasitism in birds. – Oecologia 135: 542-547.
- Swan Study Group 2005. Swan Manual. BTO, Thetford, U.K.
- Valkiūnas, G. 2005. Avian Malaria Parasites and other Haemosporidia. – CRC Press, Boca Raton, FL, USA.
- Valkiūnas, G., Anwar, A. M., Atkinson, C. T., Greiner, E. C., Paperna, I. and Peirce, M. A. 2005. What distinguishes malaria parasites from other pigmented haemosporidians? – Trends Parasitol. 21: 357-358.
- van Oers, K., Richardson, D. S., Saether, S. A. and Komdeur, J. 2010. Reduced blood parasite prevalence with age in the Seychelles Warbler: selective mortality or suppression of infection? – J. Ornithol. 151: 69-77.
- Ventim, R., Morais, J., Pardal, S., Mendes, L., Ramos, J. A. and Pérez-Tris, J. 2012. Host-parasite associations and host-specificity in haemoparasites of reed bed passerines. – Parasitol. 139: 310-316.
- Vleck, C. M., Haussmann, M. F. and Vleck, D. 2007. Avian senescence: underlying mechanisms. – J. Ornithol. 148: S611-S624.
- Waldenström, J., Bensch, S., Hasselquist, D. and Ostman, O. 2004. A new nested polymerase chain reaction method very efficient in detecting *Plasmodium* and *Haemoproteus* infections from avian blood. – J. Parasitol. 90: 191-194.
- Westerdahl, H., Waldenström, J., Hansson, B., Hasselquist, D., von Schantz, T. and Bensch, S. 2005. Associations between malaria and MHC genes in a migratory songbird. – P. Roy. Soc. B - Biol. Sci. 272: 1511-1518.
- Williams, D. D. and Williams, N. E. 1998. Aquatic insects in an estuarine environment: densities, distribution and salinity tolerance. – Freshwater Biol. 39: 411-421.
- Wilson, K. 2002. Heterogeneities in macroparasite infections: patterns and processes. In: HUDSON, P. J., RIZZOLI, A., GRENFELL, B. T., HEESTERBEEK, H. and DOBSON, A. P. (eds.) The Ecology of Wildlife Diseases - OUP, pp. 6-44.
- Wood, M. J., Cosgrove, C. L., Wilkin, T. A., Knowles, S. C. L., Day, K. P. and Sheldon, B. C. 2007. Within-population variation in prevalence and lineage distribution of avian malaria in blue tits, *Cyanistes caeruleus*. – Mol. Ecol. 16: 3263-3273.

808 Wood, S. N. 2006. Generalized Additive Models: An Introduction with R. – Chapman &
809 Hall/CRC, Boca Raton, FL, USA.
810 Woolhouse, M. E. J. and Chandiwana, S. K. 1992. A further model for temporal patterns
811 in the epidemiology of schistosome infections of snails. – Parasitol. 104: 443-449.
812 Woolhouse, M. E. J., Taylor, P., Matanhire, D. and Chandiwana, S. K. 1991. Acquired
813 immunity and epidemiology of *Schistosoma haematobium*. – Nature 351: 757-
814 759.
815 Zehtindjiev, P., Ilieva, M., Westerdahl, H., Hansson, B., Valkiunas, G. and Bensch, S.
816 2008. Dynamics of parasitemia of malaria parasites in a naturally and
817 experimentally infected migratory songbird, the great reed warbler *Acrocephalus*
818 *arundinaceus*. – Exp. Parasitol. 119: 99-110.
819